

downloaded from Stanmorephysics.com
education



Department:
Education
PROVINCE OF KWAZULU-NATAL

**NATIONAL
SENIOR CERTIFICATE**

GRADE 11

MATHEMATICS

COMMON TEST

MARCH 2019

MARKS: 75

TIME: 1½ hours

This question paper consists of 6 pages and 2 Diagram Sheets.

INSTRUCTIONS AND INFORMATION

Read the following instructions carefully before answering the questions.

1. This question paper consists of 5 questions.
2. Answer ALL the questions.
3. Clearly show ALL calculations, diagrams, graphs, etc. which you have used in determining your answers.
4. Answers only will NOT necessarily be awarded full marks.
5. You may use an approved scientific calculator (non-programmable and non-graphical), unless stated otherwise.
6. If necessary, round off answers correct to TWO decimal places, unless stated otherwise.

Diagrams are NOT necessarily drawn to scale.
7. Write neatly and legibly.
8. Diagrams for QUESTION 4.1, QUESTION 4.2, QUESTION 5.1 and QUESTION 5.2 on the DIAGRAM SHEETS provided.
9. Detach the DIAGRAM SHEETS and hand in together with your Answer Book.

QUESTION 1

1.1 Solve for x :

1.1.1 $7x^2 - 2x - 3 = 0$ (correct to TWO decimal places) (3)

1.1.2 $(x - 2)^2 - 4 = 0$ (3)

1.1.3 $\sqrt{7x + 2} + 2x = 0$ (4)

1.1.4 $x^2 - x - 56 < 0$ (3)

1.2 Solve for x and y simultaneously: $2x + y = 1$ and $2x^2 - xy + y^2 = 4$ (6)
[19]

QUESTION 2

2.1 Solve for x without the use of a calculator : $x^{\frac{3}{4}} = 64$ (2)

2.2 Simplify without the use of a calculator :

2.2.1 $\frac{5^{-x} \cdot 125^{1-x} \cdot 25^{2x}}{5}$ (3)

2.2.2 $\sqrt{12} - \sqrt{147} + 3^{1,5}$ (3)

2.3 If $\frac{5^{2006} - 5^{2004} + 24}{5^{2004} + 1} = a$, calculate a without the use of a calculator. (3)
[11]

QUESTION 3

ANSWER QUESTION 3 WITHOUT USING A CALCULATOR.

3.1 Given: $\tan \theta = -\frac{9}{40}$ and $180^\circ < \theta < 360^\circ$.
Use a sketch to determine the value of $\sin \theta + \cos \theta$. (4)

3.2 Simplify fully:

$$\frac{\sin(90^\circ - \theta) \cdot \tan(360^\circ - \theta) \cdot \sin(\theta - 180^\circ)}{1 - \cos^2 \theta} \quad (6)$$

3.3 Determine the value of the following in terms of p , if $\cos 32^\circ = p$:


3.3.1 $\cos 212^\circ$ (2)

3.3.2 $\sin(-328^\circ)$ (3)
[15]

GIVE REASONS FOR YOUR STATEMENTS AND CALCULATIONS IN QUESTIONS 4 AND 5.

QUESTION 4


4.1 In the diagram, O is the centre of the circle and R is a point on chord ST , such that OR is perpendicular to ST .



Prove the theorem which states that $SR = RT$. (5)

downloaded from Stanmorephysics.com

- 4.2 In the diagram, O is the centre of the circle and D is a point on chord AC such that $AD = DC = 24$ mm. OD is drawn and produced to meet the circle at B . $OD = 7$ mm. OA and OC are drawn.




Calculate the length of BD .

(5)
[10]

QUESTION 5

- 5.1 In the diagram two circles intersect at L and H . O is the centre of the circle passing through F , G , H and L . GO and LO are drawn. $LHJK$ is a cyclic quadrilateral. FLK and GHJ are straight lines. $\hat{GOL} = 136^\circ$



5.1.1 Calculate the size of \hat{F} .

(2)


GRADE 11-NSC

5.1.2

Calculate the size of \hat{K} .

(4)

- 5.2 In the diagram, O is the centre of the circle. Diameter BOA is produced to E such that EDG is a tangent to the circle at D . C is a point on the circle such that $BA \parallel CD$. AD , BD and AC are drawn. F is a point of intersection of AC and BD . Let $\hat{B} = x$.




- 5.2.1 Write down, with reasons, four other angles each equal to x . (6)
- 5.2.2 Determine the size of \hat{E} in terms of x . (4)
- 5.2.3 Prove that CA is a tangent to the circle passing through A , D and E . (4)
- [20]**

TOTAL: 75

NAME & SURNAME:


DIAGRAM SHEET 1

QUESTION 4.1



TEAR OFF

QUESTION 4.2




downloaded from Stanmorephysics.com


NAME & SURNAME:

DIAGRAM SHEET 2

QUESTION 5.1



QUESTION 5.2



TEAR OFF



education

Department:
Education
PROVINCE OF KWAZULU-NATAL

MATHEMATICS

COMMON TEST

MARCH 2019

MARKING GUIDELINES

**NATIONAL
SENIOR CERTIFICATE**

GRADE 11

downloaded from Stanmorephysics.com

MARKS: 75

These marking guideline consists of 8 pages.

GEOMETRY • MEETKUNDE	
S	A mark for a correct statement (A statement mark is independent of a reason)
	<i>'n Punt vir 'n korrekte bewering</i> <i>('n Punt vir 'n bewering is onafhanklik van die rede)</i>
R	A mark for the correct reason (A reason mark may only be awarded if the statement is correct)
	<i>'n Punt vir 'n korrekte rede</i> <i>('n Punt word slegs vir die rede toegeken as die bewering korrek is)</i>
S/R	Award a mark if statement AND reason are both correct
	<i>Ken 'n punt toe as die bewering EN rede beide korrek is</i>

QUESTION 1

1.1.1	$7x^2 - 2x - 3 = 0$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $x = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(7)(-3)}}{2(7)}$ $x = -0,53 \text{ or } x = 0,81$	<ul style="list-style-type: none"> ✓ substituting in correct formula ✓ x-values ✓ x-values <p style="text-align: right;">(3)</p>
1.1.2	$(x - 2)^2 - 4 = 0$ $(x - 2)^2 = 4$ $x - 2 = \pm 2$ $x = 4 \text{ or } x = 0$ <p style="text-align: center;">OR</p> $(x - 2)^2 - 4 = 0$ $x^2 - 4x + 4 - 4 = 0$ $x^2 - 4x = 0$ $x(x - 4) = 0$ $x = 4 \text{ or } x = 0$	<ul style="list-style-type: none"> ✓ isolate $(x - 2)^2$ ✓ ± 2 ✓ both answers <p style="text-align: center;">OR</p> <ul style="list-style-type: none"> ✓ $x^2 - 4x + 4$ ✓ factors ✓ both answers <p style="text-align: right;">(3)</p>
1.1.3	$\sqrt{7x + 2} + 2x = 0$ $(\sqrt{7x + 2})^2 = (-2x)^2$ $7x + 2 = 4x^2$ $4x^2 - 7x - 2 = 0$ $(4x + 1)(x - 2) = 0$ $x = -\frac{1}{4} \text{ or } x = 2$ $\therefore x = -\frac{1}{4} \text{ only}$	<ul style="list-style-type: none"> ✓ isolate $\sqrt{7x + 2}$ ✓ standard form ✓ factors ✓ correct solution <p style="text-align: right;">(4)</p>



<p>1.1.4</p>	$x^2 - x - 56 < 0$ $(x - 8)(x + 7) < 0$ <p>CV $x = 8$ or $x = -7$</p> $\begin{array}{ccccccc} & + & & - & & + & \\ & & & & & & \\ - & 7 & & 8 & & & \end{array}$ $-7 < x < 8$	<p>✓ correct factors</p> <p>✓✓ correct solution</p> <p>(3)</p>
<p>1.2</p>	$2x + y = 1 \text{ and } 2x^2 - xy + y^2 = 4$ $y = 1 - 2x$ $2x^2 - x(1 - 2x) + (1 - 2x)^2 = 4$ $2x^2 - x + 2x^2 + 1 - 4x + 4x^2 = 4$ $8x^2 - 5x - 3 = 0$ $(8x + 3)(x - 1) = 0$ $x = -\frac{3}{8} \text{ or } x = 1$ $y = 1 - 2\left(-\frac{3}{8}\right) \text{ or } y = 1 - 2(1)$ $y = 1\frac{3}{4} \text{ or } y = -1$ <p>OR</p> $2x + y = 1 \text{ and } 2x^2 - xy + y^2 = 4$ $x = \frac{1 - y}{2}$ $2\left(\frac{1 - y}{2}\right)^2 - y\left(\frac{1 - y}{2}\right) + y^2 = 4$ $2\left(\frac{1 - 2y + y^2}{4}\right) - y\left(\frac{1 - y}{2}\right) + y^2 - 4 = 0$ $1 - 2y + y^2 - y + y^2 + 2y^2 - 8 = 0$ $4y^2 - 3y - 7 = 0$ $(4y - 7)(y + 1) = 0$ $y = 1\frac{3}{4} \text{ or } y = -1$ $x = \frac{1 - 1\frac{3}{4}}{2} \text{ or } x = \frac{1 - (-1)}{2}$ $x = -\frac{3}{8} \text{ or } x = 1$	$y = 1 - 2x$ <p>✓ substitution</p> <p>✓ standard form</p> <p>✓ factors</p> <p>✓ x values</p> <p>✓ y values</p> <p>(6)</p> <p>OR</p> $x = \frac{1 - y}{2}$ <p>✓ substitution</p> <p>✓ standard form</p> <p>✓ factors</p> <p>✓ y values</p> <p>✓ x values</p> <p>(6)</p>

[19]

QUESTION 2

<p>2.1</p>	$x^{\frac{3}{4}} = 64$ $\left(x^{\frac{3}{4}}\right)^{\frac{4}{3}} = (2^6)^{\frac{4}{3}}$ $x = 256 \text{ or } 2^8$	<p>✓ raising both sides to the $\frac{4}{3}$</p> <p>✓ answer</p> <p>(2)</p>
<p>2.2.1</p>	$\frac{5^{-x} \cdot 125^{1-x} \cdot 25^{2x}}{5}$ $= \frac{5^{-x} \cdot (5^3)^{1-x} \cdot (5^2)^{2x}}{5}$ $= \frac{5^{-x} \cdot 5^{3-3x} \cdot 5^{4x}}{5}$ $= 5^{-x+3-3x+4x-1}$ $= 5^2$ $= 25$	<p>✓ rewriting as base 3</p> <p>✓ using exponential rules</p> <p>✓ answer</p> <p>(3)</p>
<p>2.2.2</p>	$\sqrt{12} - \sqrt{147} + 3^{1,5}$ $= \sqrt{4 \times 3} - \sqrt{49 \times 3} + 3^{\frac{3}{2}}$ $= 2\sqrt{3} - 7\sqrt{3} + \sqrt{9 \times 3}$ $= 2\sqrt{3} - 7\sqrt{3} + 3\sqrt{3}$ $= -2\sqrt{3}$	<p>✓ simplifying surds</p> <p>✓ $3\sqrt{3}$</p> <p>✓ answer</p> <p>(3)</p>
<p>2.3</p>	$\frac{5^{2006} - 5^{2004} + 24}{5^{2004} + 1} = a$ $\frac{5^{2004}(5^2 - 1) + 24}{5^{2004} + 1} = a$ $\frac{5^{2004}(24) + 24}{5^{2004} + 1} = a$ $\frac{24(5^{2004} + 1)}{5^{2004} + 1} = a$ $a = 24$	<p>✓ factorising</p> <p>✓ factorising</p> <p>✓ answer</p> <p>(3)</p>
<p>[11]</p>		

QUESTION 3

<p>3.1</p>	 $x^2 + y^2 = r^2$ $40^2 + (-9)^2 = r^2$ $r = 41$ $\frac{\sin \theta + \cos \theta}{41} = \frac{-9}{41} + \frac{40}{41}$ $= \frac{31}{41}$	<p>✓ correct sketch in 4th quadrant</p> <p>✓ value of r</p> <p>✓ substitution</p> <p>✓ answer</p> <p style="text-align: right;">(4)</p>
<p>3.2</p>	$\frac{\sin(90^\circ - \theta) \cdot \tan(360^\circ - \theta) \cdot \sin(\theta - 180^\circ)}{1 - \cos^2 \theta}$ $= \frac{\cos \theta \cdot -\tan \theta \cdot -\sin \theta}{\sin^2 \theta}$ $= \frac{\cos \theta \cdot -\frac{\sin \theta}{\cos \theta} \cdot -\sin \theta}{\sin^2 \theta}$ $= 1$	<p>$\cos \theta$</p> <p>$-\tan \theta$</p> <p>$-\sin \theta$</p> <p>$\sin^2 \theta$</p> <p>$\frac{\sin \theta}{\cos \theta}$</p> <p>✓ answer</p> <p style="text-align: right;">(6)</p>
<p>3.3.1</p>	$\cos 212^\circ$ $= \cos(180^\circ + 32^\circ)$ $= -\cos 32^\circ$ $= -p$	<p>$-\cos 32^\circ$</p> <p>✓ answer</p> <p style="text-align: right;">(2)</p>
<p>3.3.2</p>	$\sin(-328^\circ)$ $= \sin 32^\circ$ $= \sqrt{1 - \cos^2 32^\circ}$ $= \sqrt{1 - p^2}$ 	<p>$\sin 32^\circ$</p> <p>✓ correct sketch or identity</p> <p>✓ answer</p> <p style="text-align: right;">(3)</p>
<p>[15]</p>		

QUESTION 4

<p>4.1</p>	<div style="text-align: center;"> </div> <p>Construction: Draw OS and OT.</p> <p>Proof: In $\triangle OSR$ and $\triangle OTR$: 1. $OS = OT$ [radii] 2. $OR = OR$ [common] 3. $\hat{SRO} = \hat{TRO} = 90^\circ$ [$\angle s$ on a straight line] $\therefore \triangle OSR \equiv \triangle OTR$ [90°; H; S] $\therefore SR = RT$ [$\equiv \Delta s$]</p>	<p>✓ construction</p> <p>✓ S/R ✓ S (OR is common) ✓ S/R ✓ S/R</p> <p style="text-align: right;">(5)</p>
<p>4.2</p>	<p>$OD \perp AC$ [line from centre to midpoint of chord] $OA^2 = AD^2 + OD^2$ [Pythagoras] $= 24^2 + 7^2$ $= 625$ $OA = 25 \text{ mm}$ $OB = OA$ [radii] $\therefore BD = 25 - 7 = 18 \text{ mm}$</p>	<p>✓ S/R ✓ S/R</p> <p>✓ length of the radius ✓ S/R ✓ answer</p> <p style="text-align: right;">(5)</p>
<p>[10]</p>		

downloaded from Stanmorephysics.com

QUESTION 5

5.1.1	$\hat{F} = \frac{1}{2} \hat{GOL}$ $= 68^\circ$ <p>[∠ at centre = 2 × ∠ at circumference]</p>	✓ R ✓ answer (2)
5.1.2	$\hat{F} + \hat{H}_1 = 180^\circ$ <p>[opp. ∠ s of cyclic quadrilateral]</p> $\hat{H}_1 = 180^\circ - 68^\circ$ $= 112^\circ$ $\hat{K} = \hat{H}_1$ <p>[ext. ∠ of cyclic quadrilateral]</p> $= 112^\circ$ <p>OR</p> $\hat{H}_2 = \hat{F} = 68^\circ$ <p>[ext. ∠ of cyclic quadrilateral]</p> $\hat{K} = 112^\circ$ <p>[opp. ∠ s of cyclic quadrilateral]</p>	✓ R ✓ size of \hat{H}_1 ✓ R ✓ answer (4) OR ✓ S ✓ R ✓ S ✓ R (4)
5.2.1	$\hat{C} = x$ <p>[∠ s in the same segment]</p> $\hat{A}_1 = \hat{C} = x$ <p>[alt. ∠ s; BA CD]</p> $\hat{D}_2 = \hat{A}_1 = x$ <p>[∠ s in the same segment]</p> $\hat{D}_4 = \hat{B} = x$ <p>[tan-chord theorem]</p> <p>OR</p> $\hat{D}_2 = x$ <p>[alt. ∠ s; BA CD]</p> $\hat{A}_1 = \hat{D}_2 = x$ <p>[∠ s in the same segment]</p> $\hat{C} = \hat{B} = x$ <p>[∠ s in the same segment]</p> $\hat{D}_4 = \hat{B} = x$ <p>[tan-chord theorem]</p>	✓ S ✓ R ✓ S/R ✓ S/R ✓ S ✓ R (6) OR ✓ S/R ✓ S ✓ R ✓ S/R ✓ S ✓ R (6)
5.2.2	$\hat{D}_3 = 90^\circ$ <p>[∠ in a semicircle]</p> $\hat{E} = 180^\circ - (\hat{B} + \hat{BDE})$ <p>[sum of ∠ s in Δ]</p> $= 180^\circ - (x + 90^\circ + x)$ $= 90^\circ - 2x$ <p>OR</p> $\hat{D}_3 = 90^\circ$ <p>[∠ in a semicircle]</p> $\hat{E} + \hat{CDE} = 180^\circ$ <p>[co-interior ∠ s; BA CD]</p> $\hat{E} = 180^\circ - \hat{CDE}$ $= 180^\circ - (x + 90^\circ + x)$ $= 90^\circ - 2x$	✓ S ✓ R ✓ S ✓ answer (4) OR ✓ S ✓ R ✓ S ✓ answer (4)

<p>5.2.3</p>	<p> $\hat{A}_2 = 180^\circ - (\hat{B} + \hat{D}_3 + \hat{A}_1)$ [sum of \angles in $\triangle ABD$] $= 180^\circ - (x + 90^\circ + x)$ $= 90^\circ - 2x$ $\therefore \hat{A}_2 = \hat{E}$ [both = $90^\circ - 2x$] $\therefore AE$ is a tangent to the circle through A, D and E [converse: tan-chord-theorem] </p> <p>OR</p> <p> $\hat{D}_1 = 180^\circ - (\hat{D}_2 + \hat{D}_3 + \hat{D}_4)$ [\angles on a straight line] $= 180^\circ - (x + 90^\circ + x)$ $= 90^\circ - 2x$ $\hat{D}_1 = \hat{A}_2$ [tan-chord-theorem] $\therefore \hat{A}_2 = 90^\circ - x$ $\therefore \hat{A}_2 = \hat{E}$ [both = $90^\circ - 2x$] $\therefore AE$ is a tangent to the circle through A, D and E [converse: tan-chord-theorem] </p>	<p> \checkmark S $\hat{A}_2 = 90^\circ - 2x$ $\hat{A}_2 = \hat{E}$ \checkmark R (4) </p> <p>OR</p> <p> \checkmark S $\hat{A}_2 = 90^\circ - 2x$ $\hat{A}_2 = \hat{E}$ \checkmark R (4) </p>
[20]		

TOTAL MARKS: 75

downloaded from Stanmorephysics.com